arXiv:1209.0221v1 [math.GT] 2 Sep 2012

Pictures of the Chabauty space of $\mathbb{C}^*$

Hyungryul Baik & Lucien Clavier
Department of Mathematics
310 Malott Hall, Cornell University
Ithaca, New York 14853-4201 USA

September 4, 2012

Abstract

In this note not intended for publication, we provide pictures for the Chabauty space of $\mathbb{C}^*$.

Lemma 1 and Propositions 2 and 3 are not new, and can be found in [Bac11] for instance. This note was announced in [BC12].

Recall that any closed subgroup of $\mathbb{C}$ is isomorphic to exactly one of the following groups: $\{0\}$, $\mathbb{Z}$, $\mathbb{Z}^2$, $\mathbb{R}$, $\mathbb{Z}\times\mathbb{R}$, $\mathbb{C}$.

Lemma 1. *The followings are all the closed subgroups of* $\mathbb{C}$ *containing* $2i\pi$*.*

- $A^m := (2\pi/m)i\mathbb{Z}$ *for some* $m \in \mathbb{N}$*,*
- $B^m_z := z\mathbb{Z} + (2\pi/m)i\mathbb{Z}$ *for some* $m \in \mathbb{N}$ *and for* $z \in \mathbb{C}$ *with* $\mathrm{Re}(z) > 0$ *and* $\mathrm{Im}(z) \in [0, m]$*,*
- $C_x := x\mathbb{Z} + i\mathbb{R}$ *for* $x > 0$*,*
- $D^m_t := (2\pi/m)i\mathbb{Z} + (1 + it)\mathbb{R}$ *with* $t \in \mathbb{R}$ *and* $m \in \mathbb{N}$*.*
- $A^0 = C_\infty := i\mathbb{R}$*,*
- $C_0 := \mathbb{C}$*.*

Proof. Let Γ be a closed subgroup of $\mathbb{C}$ containing $2i\pi$. Then Γ contains A^1. If Γ is discrete, it must contain A^m for some maximal m; if Γ is isomorphic to $\mathbb{R}$, it must be $C_\infty = i\mathbb{R}$. Also, if Γ is a lattice containing A^m for a maximal m, it must be of the form B^m_z for z verifying $\mathrm{Re}(z) > 0$ and $\mathrm{Im}(z) \in [0, m]$. Thus, suppose Γ is isomorphic to $\mathbb{R}\times\mathbb{Z}$. There are two cases.
Case 1: A^m is the $\mathbb{Z}$ part. Then Γ, as a set, is the union of parallel lines of finite slope t passing through the points of A^m. Thus Γ is D^m_t.
Case 2: A^m is contained in the $\mathbb{R}$ part. Then Γ contains C_∞ and is the union of vertical lines equally spaced in the horizontal direction. Hence $\Gamma = C_x$ for some $x > 0$.

□

Proposition 2. *In the Chabauty topology, we have the following convergence results.*

- $A^{m_n} \to \begin{cases} A^0 = i\mathbb{R} \text{ if } m_n \to \infty \\ A^m \text{ if } m_n \to m \in \mathbb{N} \end{cases}$
- $B^{m_n}_{z_n} \to \begin{cases} C_x \text{ if } m_n \to \infty \text{ and } \mathrm{Re}(z_n) \to x \\ A^m \text{ if } m_n \to m \in \mathbb{N} \text{ and } \mathrm{Re}(z_n) \to \infty \\ B^m_z \text{ if } m_n \to m \in \mathbb{N} \text{ and } z_n \to z \text{ with } \mathrm{Re}(z) \in (0,\infty) \end{cases}$
- $C_{x_n} \to \begin{cases} C_0 = \mathbb{C} \text{ if } x_n \to 0 \\ C_\infty = i\mathbb{R} \text{ if } x_n \to \infty \\ C_x \text{ if } x_n \to x \in (0,\infty) \end{cases}$
- $D^{m_n}_{t_n} \to \begin{cases} \mathbb{C} \text{ if } m_n \to \infty \text{ or } t_n \to \pm\infty \\ D^m_t \text{ if } t_n \to t \in \mathbb{R} \text{ and } m_n \to m \in \mathbb{N} \end{cases}$

Proof. The proofs of these assertions are either easier than or similar to the proof of Proposition 3 below; they are therefore left to the reader to check. □

Proposition 3. *Let $(z_n = x_n + 2i\pi\theta_n)$ be a converging sequence of complex numbers, with $x_n > 0$, $x_n \to 0$, $\theta_n \in [0,1]$, $\theta_n \to \theta$. If θ is rationnal, say $\theta = p/q$ with p, q coprime positive integers, define for all n t_n to be the slope of the line passing through 0 and $qz_n - 2i\pi p$, i.e.*

$$t_n = \frac{2\pi}{x_n}(\theta_n - \theta).$$

Then the limit in the Chabauty topology of the sequence $B^m_{z_n}$ is

$$\begin{cases} D^{\mathrm{lcm}(m,q)}_t \text{ if } t_n \to t \in \mathbb{R} \\ \mathbb{C} \text{ if } t_n \to \pm\infty \end{cases}$$

If θ is irrationnal then $B^m_{z_n} \to \mathbb{C}$.

Proof. The case where θ is irrationnal is immediate; let us suppose that $\theta \in \mathbb{Q}$.

Note, just by drawing all lines of slope t_n passing through points of $B^m_{z_n}$, that $B^m_{z_n} \subset D^{\mathrm{lcm}(m,q)}_{t_n}$. Moreover, a closer look on the intersection between all those lines and the imaginary axis $i\mathbb{R}$ shows that on every line of $D^{\overline{\frac{mq}{\gcd(pm,q)}}}_{t_n}$ there is actually at least one point of $B^m_{z_n}$, as soon as θ_n is close enough to θ. Since p and q are coprime, $\frac{mq}{\gcd(pm,q)} = \mathrm{lcm}(m,q)$, thus there is at least one point of $B^m_{z_n}$ on each line of $D^{\mathrm{lcm}(m,q)}_{t_n}$.

Finally, since $x_n \to 0$, we can find for every $\epsilon > 0$ an integer N large enough so that for all $n \geq N$, $B^{2\pi/m}_{z_n}$ ϵ-fills $D^{\mathrm{lcm}(m,q)}_{t_n}$ (i.e. every point of $D^{\mathrm{lcm}(m,q)}_{t_n}$ is at distance at most ϵ of a point of $B^{2\pi/m}_{z_n}$). Therefore the Hausdorff distance between $B^{2\pi/m}_{z_n}$ and $D^{\mathrm{lcm}(m,q)}_{t_n}$ tends to zero, and we are done. □

Let us interpret these results geometrically.

First, let us describe the space of subgroups D_t^m, for $m \in \mathbb{N}$ and $t \in [-\infty, \infty]$. By Proposition 2, $D_t^m \to \mathbb{C}$ for any m if $t \to \pm\infty$. Thus we get a bouquet of circles; one circle for each $m \in \mathbb{N}$, the wedge point corresponding to the total subgroup $\mathbb{C}$. We also know that when $m \to \infty$, $D_t^m \to \mathbb{C}$ for any t. Thus when we increase m, the corresponding circle in the bouquet shrinks down to the wedge point. We call this space the D-bouquet. See Figure 1 below.


D_t^2, $t \in [-\infty, \infty]$
D_t^3, $t \in [-\infty, \infty]$
D_t^1 , $t \in [-\infty, \infty]$
D_t^m, $t \in [-\infty, \infty]$

Figure 1: The D-bouquet, a bouquet of circles; the wedge point represents the total subgroup $\mathbb{C}$.

Now set $m \in \mathbb{N}$ to be fixed. We would like to see how the closure of the space of subgroups B_z^m looks like, for z verifying $\mathrm{Re}(z) > 0$ and $\mathrm{Im}(z) \in [0, m]$.

But since two subgroups B_z^m, $B_{z'}^m$ for z, z' as above, are the same if and only if $z = z'$ mod $2i\pi/m$, the space of B_z^m's is the cylinder

$$\{z = x + iy;\ x > 0,\ y \in [0, m]\} \Big/ (x \sim x + 2i\pi/m).$$

By Proposition 2, if $x \to \infty$, then $B_z^m \to A_m$ in the Chabauty topology. Therefore the space becomes a cone in the right direction. The identification of the other end is more complicated. Say $x \to 0$ and $y \to 2\pi\theta$ with $\theta \in [0, m]$. By Proposition 3, there are two cases to consider.

Case 1: θ is irrational. Then B_z^m converges to $\mathbb{C}$.

Case 2: θ is rational, say $\theta = p/q$. Then $B_z^m \to D_t^{\mathrm{lcm}(m,q)}$ where $t = \lim \frac{2\pi}{x_n}(\theta_n - \theta)$. For p and q fixed, every possible limit for t is possible; hence we have to blow up the point $0 + 2i\pi p/q$ at the left of the cylinder to a segment corresponding to $D_t^{\mathrm{lcm}(m,q)}$ with $t \in [-\infty, \infty]$. Now since $D_{\pm\infty}^{\mathrm{lcm}(m,q)} = \mathbb{C}$, we still have to pinch the endpoints of that segment to a point, as in Figure 2 below.

Therefore the left end of the cylinder, where $x = 0$, is glued to D-bouquet, in such a way that all points $2i\pi\theta$ with θ irrational are collapsed to the wedge point of the bouquet, and the other points are blown up to some circles of the bouquet.

Note that $\theta_1 = p_1/q_1$ and $\theta_2 = p_2/q_2$ are blown up to the *same circle*, as long as $\mathrm{lcm}(m, q_1) = \mathrm{lcm}(m, q_2)$. Thus if $m = 1$, then this end is exactly the D-bouquet; but whenever $m > 1$, the end is glued to a proper subbouquet, containing only petals of index in $m\mathbb{N}$.



Figure 2: At the left of the cylinder, each point $2i\pi\theta$ with θ rational is blown-up to a circle, resulting from a pinching. First step: $2i\pi\theta$ (in red in the first picture) is blown-up to a segment (in red in the second picture). In orange, each ray represents a locus of constant slope t_n (see Proposition 3 for notations). Second step: force the end-points of this segment to get closer and closer together (third picture) until the pinching (last picture).

We call the resulting space the mth layer, noted L_m.

We can now collect every result of Propositions 2 and 3 into a global picture for the Chabauty space of $\mathbb{C}^*$ (see Figure 3).



Figure 3: The Chabauty space of $\mathbb{C}^*$

References

[Bac11] Ismaël Bachy. *Enrichissements de Siegel*. PhD thesis, Université de Provence, 2011.

[BC12] Hyungryul Baik and Lucien Clavier. The space of geometric limits of abelian subgroups of PSL$(2, \mathbb{C})$. 2012. (unpublished).